







* 4 POs of the 11 associated to Assomela

Registration number

S-P-00369

PCR

2019:01 Fruit and nuts. v 1.01

Programme

The International EPD® System www.environdec.com

Date of publication

2012/11/08

Date of validity

2028/12/17

Date of revision

2023/12/17

Version

12

CPC Code

013 Friuts and nut

Information related to

2022 harvest

Programme operator

EPD International AB

Sector EPD

This EPD has been developed in accordance with ISO 14025. An EPD should provide current information, and may be updated if conditions change. The stated validity is therefore subject to the continued registration and publication at www.environdec.com




Assomela

Assomela is the Italian association of Producer Organizations (POs), representing **70% of the national apple production**.

Assomela brings together POs VI.P (Val Venosta) VOG Marlene and VOG Products of the province of Bolzano, Melinda, La Trentina and Mezzacorona of the province of Trento, Melapiù of the region Emilia Romagna, Rivoira, Gullino and Lagnasco of the region Piemonte, Frutta Friuli SCA of the region Friuli Venezia Giulia.

The mission of Assomela is the representation of associated producers' interests by coordinating and realizing research projects of common interest.





Producers' organizations involved in the project

La Trentina (Trento)

La Trentina represents about **1,000 farms** that operate in Trentino. They are grouped together into **a single cooperative**.

The annual production is about **50,000 tons** of apples, cultivated with kiwi, cherries and plums on around **1,100** hectares

www.latrentina.it



Melinda (Cles)

The Melinda Consortium since 1989 has assembled **16** cooperatives and **3,600 production units**.

Cooperatives are base in the Non and Sole and produce about **420,000 tons** a year on around **6,700 hectares**.

www.melinda.it



VOG (Terlano)

VOG is the Consortium of fruit and vegetable Cooperatives from Alto Adige. It brings together 11 cooperatives and 4,600 producers on a surface of 11,000 hectares. Annual crop is around 530,000 tons.

www.vog.it



VI.P (Laces)


VI.P brings together **6 cooperatives** made up of **1,600 producers** from Val Venosta. On **5,300 hectares**, annual yield is about **290,000 tons** of apples.

www.vip.coop





The cultivation areas






The apple

Apples are the fruit, more precisely false fruits, of the apple tree. The plant comes from **central Asia** and is present in Italy with around **2,000 nominal varieties**. It is hard to know how many varieties there are, in particular if we consider the historical superimposition of different denominations and all the extinct or untraceable species.

Even though the natural ripening is in the period between the **beginning of August and the beginning of November**, apples are available throughout the year thanks to conservation techniques. Conservation is possible in storage facilities where apples are maintained at low temperatures and at controlled atmosphere.

When requested by the market, apples are then withdrawn from these **storage plants** and taken to the **processing plants** in which they are sized, selected and packaged.

The apples we studied are cultiveted in the area analysed in this project. Varieties can differ for agronomic aspects, such as the **quantity of necessary feritlizers** and **yield**, but differences in the environmental impacts are not significative. For this reason, the resulting impact can be considered to be referring to an "average apple". It is worth noting that the variety **Golden Delicious** is the most produced by the associated producers.





Functional unit

The data presented are referred to **1 kg of apples sold unpacked.** The apples subject of the declaration are intended for direct consumption. The shelf life varies from a few days to a few weeks, depending on the variety and on the temperature of the preservation environment. Product losses due to the distribution and sales phase were not taken into consideration.

This EPD considers average values and represents an average product, not available for purchase on the market.

Product contents

Apples contain many vitamins and minerals, especially **vitamin C** and **potassium**. They are rich in **pectin**, a food fiber very important for a good digestion and for an extended sensation of satiety.

They contain **flavonoids**, which have a positive effectv on the immune system, are anti-inflammatory and may reduce the risk in contracting some kind of cancer. Shown below the detail of apples' nutritional properties.

ENERGY VALUE	ELEMENTS	MINERALS	VITAMINS	
Energy – 53 kcal	Water – 82.5 g	Water – 82.5 g Potassium – 125 mg		
	Proteins – 0.3 g	Phosphorus – 12 mg	Vitamin B1 – 0.02 mg	
	Fat – 0.1 g	Calcium – 7 mg	Vitamin B2 – 0.02 mg	
	Carbohydrates – 13.7 g	Sodium – 2 mg	Vitamin C – 6 mg	
	Food fibers – 2 g	Iron – 0.3 mg	Niacin – 0.3 mg	



The methodology for the calculation

The data presented in this environmental declaration were calculated by analyzing the impacts of the activities carried out during all the phases from the **orchard** to the **product distribution** to the consumer. The nursery phase has not been considered since the average orchard duration may exceed 25 years and thus the impacts of this phase can be considered negligible. This aspect is coherent according to the PCR (Par. 4.3.1.1).

This EPD refers to the average values for the 2022 harvest:

Agricultural phase: 1/1/2022 - 31/10/2021ì2 **Production process phase:** 1/8/2022 - 31/07/2023

PHASES OF APPLES PRODUCTION

UPSTREAM

- > Production of raw materials used during the farming phase
- > Production of auxiliary materials used during the plant phase
- > Packaging materials production

CORE

- > Emissions related to the farming phase
- > Emissions related to the the plant and storing phase
- > Water consumption
- > Wastes production and treatment

DOWNSTREAM

- > Distribution
- > Packaging End of Life



FARMING

All the activities carried out between the end of an apple picking season and the beginning of the next one are part of this phase.

The main environmental factors concern oil use and water consumption, together with the use of pesticides and fertilizers.



PLANT

Storage

After the harvest, apples are immediately left in storage cells in controlled atmosphere, where they stay untill put on the market. The main environmental impact is the consumption of electric energy.

.....

Processing

Processing is the phase of the selection, washing and packing of the apples requested by the consumers.

Electric energy and water are the impacts of this phase.



PACKAGING

The packing phase occurs right after the processing phase. It refers to the production and the use of packaging materials.



DISTRIBUTION

The distribution happens both in Italy and abroad, and it principally consists for most in land transport by truck and for a small percentage in sea transport by ship.



Main hypotheses taken into account



CULTIVATION

Consumption of water and diesel oil were estimated from four POs according to the real consumption of the sample of farms. Data on other consumption (fertilizers and pesticides) have been obtained by the production specifications of the areas interested and then validated with specific information. The data about yields were evaluated according to the average age of the crops and the production volume. The data about produced waste were collected by APOT, the fruit and vegetable organization of Trentino. The land use change was not included in the calculation since almost all the orchards are in the areas subject of the study for over 20 years.



PLANT

Storage

The electric energy consumption has been estimated by dividing the total storages' consumption by the quantity of stored apples. The average value has been calculated as described in the section referring to the calculation of the means.

Processing

In this phase electric energy consumptions, water consumption and waste production have been considered. The data gathered from a sample plants were elaborated as indicated.



PACKAGING END OF LIFE

The end of life of packging primary was assessed as an average scenario of waste disposal compostable in Italy,, considered representative for the entire international context.



USAGE

It is supposed that domestic preservation happens at room temperature.
Scraps due to the possible non edible part were not considered.



DISTRIBUTION

The impacts referred to the transportation phase have been calculated by supposing a transport of about 750 km by truck and 840 km by ship, because in addition to the Italian and European markets, the distribution also concerned overseas, Asian and North African markets.



PACKAGING

The presented data refer to the selling of unpacked apples and considering the use of one biodegradable and compostable bag for 1 kg of apples. However, other packaging possibilities are presented.



The environmental impacts

ENVIRONMENTAL IMPACT INDICATORS data relating to 1 kg of product		MEASURE	UPSTREAM		CORE		DOWNSTREAM		
			Agricultural raw materials production	Packaging e auxiliary materials production	Field phase	Plant and storage	Distribution	Packaging End of life	TOTAL
	fossil	kg CO ₂ eq	5,13E-03	9,48E-03	4,20E-02	2,82E-02	6,64E-02	9,63E-05	1,51E-01
Global Warming	biogenic	kg CO₂ eq	2,86E-06	1,99E-05	1,89E-06	2,91E-03	3,79E-06	1,42E-03	4,35E-03
Potential - (GWP)	land use and land use change	kg CO ₂ eq	4,77E-06	2,07E-04	1,32E-06	2,45E-06	1,36E-06	7,42E-10	2,17E-04
	TOTAL	kg CO ₂ eq	5,14E-03	9,71E-03	4,20E-02	3,11E-02	6,64E-02	1,51E-03	1,56E-01
Acidificatio	on potential (AP)	mol H+ eq	5,29E-05	5,89E-05	4,11E-04	8,81E-05	4,32E-04	4,28E-07	1,04E-03
	n potential (EP), uatic freshwater	kg P eq	4,75E-07	7,50E-07	1,35E-05	4,34E-07	5,05E-08	1,04E-08	1,52E-05
Eutrophicatio	n potential (EP), aquatic marine	kg N eq	4,41E-06	1,47E-05	3,67E-04	1,80E-05	1,34E-04	2,51E-06	5,41E-04
Eutrophicatio	n potential (EP), terrestrial	mol N eq	9,67E-05	1,36E-04	2,07E-03	1,98E-04	1,46E-03	1,64E-06	3,96E-03
Photochemical ozone creation potential (POCP)		kg NMVOC eq	1,53E-05	4,47E-05	4,98E-04	8,77E-05	4,61E-04	9,20E-07	1,11E-03
Ozone depletion potential (ODP)		kg CFC11 eq	1,67E-08	4,40E-10	5,82E-10	7,58E-10	1,37E-09	3,12E-12	1,98E-08
Abiotic depletion potential (ADP) for minerals and metals		kg Sb eq	2,39E-09	3,17E-09	1,26E-09	7,39E-10	2,07E-09	3,37E-13	9,64E-09
Abiotic depletion	potential (ADP) r fossil resources	МЈ	6,93E-02	2,48E-01	4,84E-01	3,87E-01	8,71E-01	5,14E-04	2,06E+00
Water deprivation potential (WDP)		m3 depriv.	3,06E-03	1,12E-02	2,27E+00	6,25E-01	7,81E-04	9,80E-03	2,92E+00

The values reported are the result of rounding. That is why the results might be slightly different from the sum of the individual components.



The environmental impacts

USE OF RESOURCES data relating to 1 kg of product			UPSTREAM		CORE		DOWNSTREAM		
		MEASURE	Produzione materie prime agricole	Packaging e materiali ausiliari di stabilimento	Fase di campo	Stabilimento e stoccaggio	Distribuzione	Fine vita packaging	TOTAL
Renewable energy resources	Use as energy carrier	MJ, net calorific value	2,05E-03	2,30E-02	7,93E-04	1,55E+00	2,20E-03	3,94E-05	1,57E+00
	Use as raw materials	MJ, net calorific value	0,00E+00	3,41E-02	0,00E+00	0,00E+00	0,00E+00	0,00E+00	3,41E-02
	TOTAL	MJ, net calorific value	2,05E-03	5,70E-02	7,93E-04	1,55E+00	2,20E-03	3,94E-05	1,61E+00
Non renewable energy resources _	Use as energy carrier	MJ, net calorific value	7,77E-02	1,38E-01	4,89E-01	4,52E-01	8,80E-01	5,68E-04	2,04E+00
	Use as raw materials	MJ, net calorific value	0,00E+00	1,44E-01	0,00E+00	0,00E+00	0,00E+00	0,00E+00	1,44E-01
	TOTAL	MJ, net calorific value	7,77E-02	2,81E-01	4,89E-01	4,52E-01	8,80E-01	5,68E-04	2,18E+00

The values reported are the result of rounding. That is why the results might be slightly different from the sum of the individual components.




The influence of primary packaging

The fossil GWP of 1 kg of apples with three different kinds of packaging has been calculated:

- > **Unpacked**, relating to the selling of loose apples at supermarkets (the use of one biodegradable and compostable bag made per 1 kg of apples has been taken into consideratio);
- > A 4 apples pack, relating to a cardboard tray of 4 apples with a plastic film in PVC;
- > 6 kg tray, relating to a cardboard tray with an R-PET layer containing 6 kg of apples.

For all three packaging, only the primary packaging and an average end-of-life scenario representative of the Italian situation were considered.





Information and contacts

REFERENCE							
Assomela, as EPD owner, has the sole owership, liability and responsabbility of this EPD.							
PROGRAM OPERATOR: EPD International AB, Box 210 60, SE-100 31 Stockholm, Sweden info@environdec.com							
THIRD PARTY EPD VERIFICATION							
Product category rules (PCR) Fruits and nuts 2019:01 Version 1.01 CPC code: 013 fruits and nuts	PCR review was conducted by: The Technical Committee of the International EPD® System. Chair: Filippo Sessa. Contact via info@environdec.com						
Independent verification of the declaration and data, according to ISO 14025: EPD process verification EPD verification - Third party verifier	Procedure for follow-up of data during EPD validity involves third part verifier: ✓ Yes No						
Third party verifier: Elia Rillo – rillo@studiofieschi.it www.studiofieschi.it							
Approved by: "The International EPD® System Technical Committee, supported by Secretariat							
EPDs within the same product category but from different programmes may not be comparable							
CONTACTS							
Assomela, via del Brennero 322, 28121 - Trento (TN), Italy info@assomela.it ww	ww.assomela.it Sassociazione italiana produttori di mele						
Technical support and grafic design: Life Cycle Engineering srl – Italy www.lce	engineering.eu (IE)						



Glossary

CARBON FOOTPRINT

A product carbon footprint is the total amount of greenhouse gases produced along the entire life cycle. It is expressed in equivalent mass of carbon dioxide $(CO_a-ea).$ In agriculture a significant contribution is given by the emission of nitrous oxide (N_aO) due to the fertilizers use.

www.ipcc.ch

WATER DEPRIVATION POTENTIAL

Water deprivation measures the available water remaining per unit of surface in a given watershed relative to the world average. after human and aquatic ecosystem demands have been met. This method builds on the assumption that the potential to deprive another user of water is directly proportional to the amount of water consumed and inversely proportional to the available water remaining per unit of surface and time in a region (watershed).

ACIDIFICATION POTENTIAL

It is a phenomenon for which precipitation is unusually acidic, meaning that it has substandard levels of pH. It can have harmful effects on plants, aquatic animals and infrastructure. Acid rain is caused by emissions of SO₂. NO_x and NH₃.

EUTROPHICATION POTENTIAL

It is an abnormal proliferation of vegetation in the aquatic ecosystems caused by the addition of nutrients into rivers. lakes or marine water, which determinates a lack of oxygen. The eutrophication potential is mainly influenced by emission into water of phosphates and nitrates.

PHOTOCEMICAL OZONE CREATION POTENTIAL

Production of compounds that, under the light effect, are able to promote an oxidation reaction leading to ozone production in the troposphere. The indicator is mainly influenced by VOCs (Volatile organic compounds).

OZONE DEPLETION POTENTIAL

Degradation of the stratospheric layer of the ozone involved in blocking the UV component of sunrays. Depletion is due to particularly reactive components that originate from chlorofluorocarbon (CFC) or chlorofluoromethanes (CFM). The substance employed as benchmark measure for OPD is trichlorofluoromethane. or CFC-11.

www.wulca-waterlca.org



Notes and references

Differences compared to the previous version

- The division of contributions to the upstream and core phases has been modified in line with what is reported in the Fruit and nuts PCR.
- The modeling of fertilizers was conducted following the indications in the Arable and vegetable crops PCR. The modeling of pesticides was conducted using an average dataset constructed with primary data from POs.
- The transports relating to the raw materials used in the cultivation phase (fertilizers, pesticides) and plant (auxiliary materials) were counted.
- · The tertiary packaging relating to the product sold in bulk was modelled.
- · The national residual mix has been updated to the year 2022
- For OPs VI.P and VOG, the energy mix used in 2023 has changed from GO to residual mix
- The packaging end-of-life scenarios have been updated to the year 2022
- · The characterization factors of the calculation method used have been updated to the latest version available

References

- · International EPD® System; General Programme Instructions (EPD); ver 4.0 of 2021/03/29
- · PCR 2019:01 Fruit and nuts, ver 1.01 of 2019-08-18
- · PCR 2020:07 Arable and vegetables crops, ver 1.01 of 2023-03-16
- · CPC code 013 Fruits and nuts
- · LCA applicata alla filiera di coltivazione e di distribuzione delle mele da parte degli associati in Assomela, ver 02 of 2023/12/14

The results are calculated following Version 2.0 of the list of environmental performance indicators and on the characterization factors of the EF 3.1 reference package.



The calculation of the average

The informations presented refer to **four of the eleven Producer Organizations** associated to Assomela, operating in the regions of Trentino Alto Adige, Piemonte, Lombardia and Veneto. Since the goal of this declaration is to provide information typical of the whole association, the data have been processed in a way that allows to create different averages between the organizations that participate to the project, using weight factors calculated on the basis of production volumes. In detail, the average has been organized in **three different levels**:

- (M1) the average between the information relating to all the productive units (UP) referring to a single productive organization (PO) and relating to a specific variety so that it is computable the impact of a single variety produced by a PO. In this level, the data are considered "metadata", and are not subjects of communication;
- (M2) the average between the information calculated by single PO and referred to the same variety. This value, calculated by using the single variety production quantity as weighing element, allows to estimate the average impacts of every single variety;
- (M3) the total amount of apples produced per variety constitutes the weigh used for calculating the overall average value assigned to Assomela.

AVERAGE	REFERENCE PRODUCT	"OWNER"	DESCRIPTION	COMMUNICATION
M1	Variety	OP	Impact of the variety produced by a single PO	X
M2	Variety	Assomela	Impact of the variety produced by the Association	X
M3	Average apple	Assomela	Impact of the apple produced by the Association	✓